



# Poultry Guard<sup>®</sup> Dry Litter Amendment

Poultry Guard<sup>®</sup> Dry is an acidified clay litter amendment that effectively controls ammonia in poultry houses. Poultry Guard<sup>®</sup> Dry is custom applied by professional applicators just prior to bird placement for immediate and extended ammonia control.



## Recommended Application Rates



### SUMMER

75-100 lbs  
PER 1,000 SQ FT



### WINTER

100-150 lbs  
PER 1,000 SQ FT

**Note:** use the higher rate if windrowing or applying to disease challenged flocks.



✉ [info@poultryguard.com](mailto:info@poultryguard.com)

🌐 [poultryguard.com](http://poultryguard.com)



## POULTRY GUARD® DRY LITTER AMENDMENT

# Pre-application Procedures

### 1

Close house immediately upon bird removal to retain litter temperature and drive off ammonia and moisture.

### 2

Run minimum ventilation to remove ammonia and moisture from house.

### 3

Manage litter according to company guidelines (decaking, windrowing) within 48 hours of bird removal.

### 4

**DO NOT** overwork litter to powder-like consistency.

### 5

Litter should set undisturbed for at least three days prior to application.

### 6

Set-up for chick placement and preheat per company guidelines.

### 7

Schedule Poultry Guard® Dry application as close to placement as possible.

### 8

Exhaust ammonia from house by running two tunnel fans upon applicator arrival.

# Post-application Procedures

### 1

Close house immediately.

### 2

Run minimum ventilation per company guidelines.

